

Index to Volume 78

Author Index

Page numbers of errata are in *italic*.

- Abécassis, J. *See* J. A. Ingebrecht, 721
 —. *See* F. Mabil, 231
- Aboubacar, A., J. D. Axtell, C.-P. Huang, and B. R. Hamaker. A rapid protein digestibility assay for identifying highly digestible sorghum lines, 160
- Albrecht, J. A. *See* E. S. Osseyi, 375
- Alho-Lehto, P. *See* A. Lapvetäinen, 322
- Åman, P. *See* A. A. M. Anderson, 507
- Ames, N. *See* A. S. Wesley, 69
- Andersson, A. A. M., R. Andersson, and P. Åman. Starch and by-products from a laboratory-scale barley starch isolation procedure, 507
- Andersson, R. *See* A. A. M. Anderson, 507
- Ando, H. *See* H. Tang, 111
- Antes, S., and H. Wieser. Effects of high and low molecular weight glutenin subunits on rheological dough properties and breadmaking quality of wheat, 157
 —, and —. Reoxidation behavior of wheat and rye glutenin subunits, 8
- Arendt, E. K. *See* S. Kenny, 458
- Aussenac, T., J.-L. Carceller, and D. Kleiber. Changes in SDS solubility of glutenin polymers during dough mixing and resting, 39
- Autio, K., L. Flander, A. Kinnunen, and R. Heinonen. Bread quality relationship with rheological measurements of wheat flour dough, 654
 —. *See* M. Salmenkallio-Marttila, 429
 —. *See* E. Vesterinen, 442
- Auty, M. *See* S. Kenny, 458
- Axtell, J. D. *See* A. Aboubacar, 160
- Babatunde Obilana, A. *See* T. Beta, 583
- Bagorogozo, K. *See* M. Okot-Kotber, 514
- Baik, B.-K. *See* M.-R. Lee, 701
- Bamforth, C. W. *See* M. Kanauchi, 121
- Barbeau, W. E. *See* S. S. Schwarzlaff, 93
- Barton, F. E., II. *See* E. T. Champagne, 358
 —. *See* D. S. Himmelsbach, 488
- Bean, S. R., C. Hicks, M. Tuinstra, and G. L. Lookhart. Use of SDS to extract sorghum and maize proins for free zone capillary electrophoresis (FZCE) analysis, 84
 —, and G. L. Lookhart. Factors influencing the characterization of gluten proteins by size-exclusion chromatography and multiangle laser light scattering (SEC-MALLS), 608
 —, and —. Optimizing quantitative reproducibility in high-performance capillary electrophoresis (HPCE) separations of cereal proteins, 530
 —. *See* C. Hicks, 412
- Beasley, H. L., C. L. Blanchard, and F. Bekes. Preparative method for *in vitro* production of functional polymers from glutenin subunits of wheat, 464
- Becker, A., S. E. Hill, and J. R. Mitchell. Milling—A further parameter affecting the Rapid Visco Analyser (RVA) profile, 166
- Behall, K. M. *See* B. W. Li, 387
- Bekes, F. *See* H. L. Beasley, 464
 —. *See* S. Uthayakumaran, 138
- BeMiller, J. N. *See* J. A. Gray, 236
 —. *See* K. C. Huber, 173
- Bergman, C. J., J. T. Delgado, A. M. McClung, and R. G. Fjellstrom. An improved method for using a microsatellite in the rice waxy gene to determine amylose class, 57
- Bergman, E.-L., K. Fredlund, P. Reinikainen, and A.-S. Sandberg. Development of predictive models for optimization of phytate degradation in wheat and rye during hydrothermal processing, 144
- Beta, T., A. Babatunde Obilana, and H. Corke. Genetic diversity in properties of starch from Zimbabwean sorghum landraces, 583
 —, and H. Corke. Noodle quality as related to sorghum starch properties, 417
 —, —, J. R. N. Taylor, and L. W. Rooney. NOTE: Effect of steeping treatment on pasting and thermal properties of sorghum starches, 303
- Betschart, A. A. *See* T. S. Kahlon, 481
 —. *See* E. T. Champagne, 358
- Bett-Garber, K. L., E. T. Champagne, A. M. McClung, K. A. Moldenhauer, S. D. Linscombe, and K. S. McKenzie. Categorizing rice cultivars based on cluster analysis of amylose content, protein content, and sensory attributes, 551
 —. *See* E. T. Champagne, 358
- Bietz, J. A. *See* J. Xu, 181
- Blackwell, E. L. *See* B. W. Li, 387
- Blanchard, C. L. *See* H. L. Beasley, 464
- Bloch, H. A., H. F. Darlington, and P. R. Shewry. *In vitro* binding of puroindolines to wheat starch granules, 74
- Blundell, M. J. *See* M. A. K. Partridge, 294
- Borras, F. *See* K. Seetharaman, 379
- Boykin, D. *See* R. J. Bryant, 131
- Brannan, G. L., C. S. Setser, K. E. Kemp, P. A. Seib, and K. Roozeboom. Sensory characteristics of grain sorghum hybrids with potential for use in human food, 693
- Brinck, O. *See* M. Mikola, 55
- Brites, C., and J. M. Carrillo. Influence of high molecular weight (HMW) and low molecular weight (LMW) glutenin subunits controlled by *Glu-1* and *Glu-3* loci on durum wheat quality, 59
- Brown, D. *See* A. S. Wesley, 69
- Brumovsky, J. O., and D. B. Thompson. Production of boiling-stable granular resistant starch by partial acid hydrolysis and hydrothermal treatments of high-amylose maize starch, 680
- Bryant, R. J., R. S. Kadan, E. T. Champagne, B. T. Vinyard, and D. Boykin. Functional and digestive characteristics of extruded rice flour, 131
- Büchmann, N. B., H. Josefsson, and I. A. Cowe. Performance of European artificial neural network (ANN) calibrations for moisture and protein in cereals using the Danish near-infrared transmission (NIT) network, 572
- Budde, A. D. *See* D. M. Peterson, 349
- Buendia, M. O. *See* D. Sahai, 14, 116
- Bugusu, B. A., O. Campanella, and B. R. Hamaker. Improvement of sorghum-wheat composite dough rheological properties and breadmaking quality through zein addition, 31
- Buriak, P. *See* S. K. Mehra, 210
 —. *See* Y. Tian, 578
- Busch, K. G. *See* F. Meuser, 50
- Caley, M. S. *See* O. K. Chung, 493
- Campanella, O. *See* B. A. Bugusu, 31
- Campbell, G. M., R. Herrero-Sanchez, R. Payo-Rodriguez, and M. L. Merchant. Measurement of dynamic dough density and effect of surfactants and flour type on aeration during mixing and gas retention during proofing, 272
- Cao, T. K. *See* G. H. Robertson, 538
- Carceller, J.-L. *See* T. Aussenac, 39
- Carriere, C. J. *See* G. E. Inglett, 471
 —. *See* J. Xu, 181
- Carrillo, J. M. *See* C. Brites, 59
- Carson, L., and X. S. Sun. Creep-recovery of bread and correlation to sensory measurements of textural attributes, 101
- Carvalho, C. W. P., and J. R. Mitchell. Effect of sucrose on starch conversion and glass transition of nonexpanded maize and wheat extrudates, 342
- Cerletti, P. *See* E. Sironi, 476
- Champagne, E. T., K. L. Bett-Garber, C. C. Grimm, A. M. McClung, K. A. Moldenhauer, S. Linscombe, K. S. McKenzie, and F. E. Barton II. Near-infrared reflectance analysis for prediction of cooked rice texture, 358
 —. *See* K. L. Bett-Garber, 551
 —. *See* R. J. Bryant, 131
 —. *See* D. S. Himmelsbach, 488
- Chen, C.-M., and A.-I. Yeh. Effect of amylose content on expansion of extruded rice pellet, 261
- Cheryan, M. *See* K. D. Rane, 400
- Cho, S. C. *See* J. K. Jang, 186
- Chong, C. K. *See* D. L. Phillips, 629
- Chow, F. I. *See* T. S. Kahlon, 481
- Chung, D. S. *See* D. Wang, 25
- Chung, O. K., J. B. Ohm, M. S. Caley, and B. W. Seabourn. Prediction of baking characteristics of hard winter wheat flours using computer-analyzed mixograph parameters, 493
- Colucci, F. *See* P. Novaro, 217
- Conde-Petit, B. *See* S. Hug-Iten, 421
- Cooke, P. H. *See* V. Singh, 436
- Corke, H. *See* T. Beta, 303, 417, 583

- . See D. L. Phillips, 629
 —. See Y. Tan, 666
 Courtin, C. M., G. G. Gelders, and J. A. Delcour. Use of two endoxylanases with different substrate selectivity for understanding arabinoxylan functionality in wheat flour breadmaking, 564
 Cowe, I. A. See N. B. Büchmann, 572
 D'Appolonia, B. L. See S. V. Erazo-Castrejón, 243
 Darlington, H. F. See H. A. Bloch, 74
 Debyser, W. See H. Goesaert, 453
 D'Egidio, M. G. See P. Novaro, 217
 de la Luz Martínez, J. See R. Martínez, 107
 Delcour, J. A. See C. M. Courtin, 564
 —. See H. Goesaert, 453
 —. See J. A. Ingelbrecht, 721
 Delgado, J. T. See C. J. Bergman, 257
 Delwiche, S. R. See B. G. Osborne, 142
 Dillahunty, A. L., T. J. Siebenmorgen, and A. Mauromoustakos. Effect of temperature, exposure duration, and moisture content on color and viscosity of rice, 559
 Dimberg, L. H., K. Sunnerheim, B. Sundberg, and K. Walsh. Stability of oat avenanthramides, 278
 Djuric, V. See N. Mladenov, 363
 Doehlert, D. C., and M. S. McMullen. Optimizing conditions for experimental oat dehulling, 675
 —. See C. M. Courtin, 564
 —. See S. V. Erazo-Castrejón, 243
 —. See L. A. Grant, 590
 —. See J. A. Ingelbrecht, 721
 Donald, A. M. Review: Plasticization and self assembly in the starch granule, 307
 Donelson, J. R., C. S. Gaines, T. S. Donelson, and P. L. Finney. Detection of wheat preharvest sprouting using a pregelatinized starch substrate and centrifugation, 282
 Donelson, T. S. See J. R. Donelson, 282
 Doner, L. W., and D. B. Johnston. Isolation and characterization of cellulose/arabinoxylan residual mixtures from corn fiber gum processes, 200
 Dowell, F. E. See D. Wang, 25
 Eckhoff, S. R. See S. K. Mehra, 210, 222
 —. See V. Singh, 46
 —. See Y. Tian, 578
 Elias, E. M. See L. A. Grant, 590
 Eliasson, A.-C. See A. Repeckiene, 193
 Erazo-Castrejón, S. V., D. C. Doehlert, and B. L. D'Appolonia. Application of oat oil in breadbaking, 243
 Escher, F. See S. Hug-Iten, 421
 Felker, F. C. See J. Xu, 181
 Ferrer, M. See K. Seetharaman, 379
 Finney, P. L. Effects of falling number sample weight on prediction of α -amylase activity, 485
 —. See J. R. Donelson, 282
 Fjellstrom, R. G. See C. J. Bergman, 257
 Flander, L. See K. Autio, 654
 Flores, R. A. See Y. Mao, 286
 Fredlund, K. See E.-L. Bergman, 144
 Fuhrmeister, H. See F. Meuser, 50
 Gaines, C. S. See J. R. Donelson, 282
 Gebruers, K. See H. Goesaert, 453
 Gelders, G. G. See C. M. Courtin, 564
 Gélinas, P. See M. Mercier, 36
 Gianibelli, M. C., O. R. Larroque, F. MacRitchie, and C. W. Wrigley. REVIEW: Biochemical, genetic, and molecular characterization of wheat glutenin and its component subunit, 635
 —. See S. Uthayakumaran, 138
 Goesaert, H., W. Debyser, K. Gebruers, P. Proost, J. Van Damme, and J. A. Delcour. Purification and partial characterization of an endoxylanase inhibitor from barley, 453
 Gómez-Aldapa, C. A. See J. F. Toro-Vazquez, 543
 Gordon, M. H. See E. Papantoniou, 663
 Grant, L. A., N. Vignaux, D. C. Doehlert, M. S. McMullen, E. M. Elias, and S. Kianian. Starch characteristics of waxy and nonwaxy tetraploid (*Triticum turgidum*) L. var. *durum* wheats, 590
 —. See P. J. Lent, 619
 Gray, J. A., and J. N. BeMiller. Accessibility of starch granules to fatty acid amides, 236
 Griffey, C. A. See S. S. Schwarzlaff, 93
 Griffin, V. K. See A. A. Perdon, 205
 Gril, J. See F. Mabilley, 231
 Grimm, C. C. See E. T. Champagne, 358
 Guerrieri, N. See E. Sironi, 476
 Gunasekaran, S. See C. H. Hwang, 88
 Gunnarsson, E. See A. Repeckiene, 193
 Gupta, D. K. See S. K. Mehra, 210
 Hamaker, B. R. See A. Aboubacar, 160
 —. See B. A. Bugusu, 31
 Hammond, E. W. See E. Papantoniou, 663
 Haque, E. See M. C. Pasikatan, 730
 Hareland, G. A. See F. A. Manthey, 368
 Heinio, R.-L., K.-M. Oksman-Caldentey, K. Latva-Kala, P. Lehtinen, and K. Poutanen. Effect of drying treatment conditions on sensory profile of germinated oat, 707
 Heinonen, R. See K. Autio, 654
 Henson, C. A. See D. M. Peterson, 349
 Herald, T. J. See E. Obuz, 97
 Herrero-Sanchez, R. See G. M. Campbell, 272
 Hicks, C., S. R. Bean, G. L. Lookhart, J. F. Pedersen, K. D. Kofoid, and M. R. Tuinstra. Genetic analysis of kafirins and their phenotypic correlations with feed quality traits, in vitro digestibility, and seed weight in grain sorghum, 412
 —. See S. R. Bean, 84
 Hicks, K. B. See V. Singh, 46
 Hill, A. S. See M. A. K. Partridge, 294
 Hill, S. E. See A. Becker, 166
 Himmelsbach, D. S., F. E. Barton, II, A. M. McClung, and E. T. Champagne. Protein and apparent amylose contents of milled rice by NIR-FT/Raman spectroscopy, 488
 Hofer, J. L. See T. S. Kahlon, 481
 Hristov, N. See N. Mladenov, 363
 Hsiao, S. L. K., R. Kieffer, and F. J. Zeller. Significance of *Aegilops tauschii* glutenin genes on breadmaking properties of wheat, 521
 Huang, C.-P. See A. Aboubacar, 160
 Huber, K. C., and J. N. BeMiller. Location of sites of reaction within starch granules, 173
 Hug-Iten, S., F. Escher, and B. Conde-Petit. Structural properties of starch in bread and bread model systems: Influence of an antistaling α -amylase, 421
 Hurburgh, C. R. See S. K. Singh, 330, 336
 Hwang, C. H., and S. Gunasekaran. Determining wheat dough mixing characteristics from power consumption profile of a conventional mixer, 88
 Ingelbrecht, J. A., K. Moers, J. Abécassis, X. Rouau, and J. A. Delcour. Influence of arabinoxylans and endoxylanases on pasta processing and quality: I. Production of high-quality pasta with increased levels of soluble fiber, 721
 Inglett, G. E., and C. J. Carriere. Cellulosic fiber gels prepared from cell walls of maize hulls, 471
 Jackson, D. S. See D. Sahai, 14, 116
 Jackson, R. See B. G. Osborne, 142
 Jane, J.-L. See C. Perera, 249
 —. See S. K. Singh, 315
 Jang, J. K., S. H. Lee, S. C. Cho, and Y. R. Pyun. Effect of sucrose on glass transition, gelatinization, and retrogradation of wheat starch, 186
 Johansson, E., M. L. Prieto-Linde, and J. Ö. Jönsson. Effects of wheat cultivar and nitrogen application on storage protein composition and breadmaking quality, 19
 Johnson, E. R. See A. A. Perdon, 205
 Johnson, J. M. See S. S. Schwarzlaff, 93
 Johnson, L. A. See S. K. Singh, 315, 330, 336
 Johnston, D. B., and V. Singh. Use of proteases to reduce steep time and SO₂ requirements in a corn wet-milling process, 405
 —. See L. W. Doner, 300
 Jones, B.-L. See M. Mikola, 55
 —. See D. M. Peterson, 349
 Jönsson, J. Ö. See E. Johansson, 19
 Josefsson, H. See N. B. Büchmann, 572
 Juliano, B. O. See V. M. F. Lai, 596
 Juodeikiene, G., See A. Repeckiene, 193
 Kadan, R. S. See R. J. Bryant, 131
 Kahlon, T. S., F. I. Chow, J. L. Hofer, and A. A. Betschart. Effect of wheat bran fiber and bran particle size on fat and fiber digestibility and gastrointestinal trace measurements in the rat, 481
 Kaitaranta, J. See A. Lapveteläinen, 322
 Kallio, H. See A. Lapveteläinen, 322
 Kanauchi, M., and C. W. Bamforth. Release of β -glucan from cell walls of starchy endosperm of barley, 121

- Karakatsanis, A. See M. Liakopoulou-Kyriakides, 603
- Katajisto, J. See A. Lapveteläinen, 322
- Katina, K. See M. Salmenkallio-Marttila, 429
- Kaukovirta-Norja, A. See A. Wilhelmson, 715
- Kemp, K. E. See G. L. Brannan, 693
- Kenny, S., K. Wehrle, M. Auty, and E. K. Arendt. Influence of sodium caseinate and whey protein on baking properties and rheology of frozen dough, 458
- Khan, K. See J. Zhu, 125, 737
- Kianian, S. See L. A. Grant, 590
- Kieffer, R. See S. L. K. Hsam, 521
- Kim, H. Y. L., H. W. Yeom, H. S. Lim, and S.-T. Lim. Replacement of shortening in yellow layer cakes by corn dextrins, 267
- Kim, K. O. See J. K. Park, 151
- Kim, S. S. See J. K. Park, 151
- Kinnunen, A. See K. Autio, 654
- Kleiber, D. See T. Aussenac, 39
- Kofoid, K. D. See C. Hicks, 412
- Kokini, J. L. See I. A. Lambert, 226
- Köpsel, H., Sivri, D., Ng, P. K. W., and J. F. Steffe. Effects of transglutaminase enzyme on fundamental rheological properties of sound and bug-damaged wheat flour doughs, 26
- Lafiandra, D. See F. MacRitchie, 501
- Lai, V. M. F., M.-C. Shen, A.-I. Yeh, B. O. Juliano, and C. Lii. Molecular and gelatinization properties of rice starches from IR24 and Sinandomeng cultivars, 596
- Laitila, A. See A. Wilhelmson, 715
- Lam, H. S., and A. Proctor. Note: Rapid methods for milled rice surface total lipid and free fatty acid determination, 498
- Lambert, I. A., and J. L. Kokini. Effect of L-cysteine on the rheological properties of wheat flour, 226
- Lapveteläinen, A., P. Alho-Lehto, L. Sinn, T. Laukkanen, T. Lindman, H. Kallio, J. Kaitaranta, and J. Katajisto. Relationships of selected physical, chemical, and sensory parameters in oat grain, rolled oats, and cooked oatmeal—A three-year study with eight cultivars, 322
- Larroque, O. R. See M. C. Gianibelli, 635
- Latva-Kala, K. See R.-L. Heiniö, 707
- Laukkanen, T. See A. Lapveteläinen, 322
- Lee, L., P. K. W. Ng, J. H. Whallon, and J. F. Steffe. Relationship between rheological properties and microstructural characteristics of nondeveloped, partially developed, and developed doughs, 447
- Lee, M.-R., B. G. Swanson, and B.-K. Baik. Influence of amylose content on properties of wheat starch and breadmaking quality of starch and gluten blends, 701
- Lee, S. H. See J. K. Jang, 186
- Lehtinen, P. See R.-L. Heiniö, 707
- Lent, P. J., and L. A. Grant. Effects of additives and storage temperature on staling properties of bagels, 619
- Li, B. W., E. L. Blackwell, K. M. Behall, and H. G. M. Liljeberg Elmståhl. Resistant starch and total dietary fiber content of oatrim muffins with different levels of amylose, amylopectin, and β -glucan, 387
- Liakopoulou-Kyriakides, M., A. Karakatsanis, M. Stamataoudis, and S. Psomas. Synergistic hydrolysis of crude corn starch by α -amylases and glucoamylases of various origins, 603
- Liavoga, A. See M. Okot-Kotber, 514
- Lii, C. See V. M. F. Lai, 596
- Lii, C. See Y. Lin, 647
- Liljeberg Elmståhl, H. G. M. See B. W. Li, 387
- Lim, H. S. See H. Y. L. Kim, 267
- Lim, S.-T. See H. Y. L. Kim, 267
- Lin, Y., A.-I. Yeh, and C. Lii. Correlation between starch retrogradation and water mobility as determined by differential scanning calorimetry and NMR, 647
- Lindman, T. See A. Lapveteläinen, 322
- Linscombe, S. D. See K. L. Bett-Garber, 551
- See E. T. Champagne, 358
- Lookhart, G. L. See S. R. Bean, 84, 530, 608
- See C. Hicks, 412
- Lu, Z. See C. Perera, 249
- Lukow, O. M. See A. S. Wesley, 69
- Mabille, F., J. Gril, and J. Abecassis. Mechanical properties of wheat seed coats, 231
- MacRitchie, F., and D. Lafiandra. Review: Use of near-isogenic wheat lines to determine protein composition-functionality relationships, 501
- See M. C. Gianibelli, 635
- See H. Singh, 526
- Manthey, F. A., and G. A. Hareland. Effects of break-roll differential on semolina and spaghetti quality, 368
- Mao, Y., and R. A. Flores. Mechanical starch damage effects on wheat flour tortilla texture, 286
- See M. D. Pearce, 354
- Martínez, J. de la Luz. See R. Martínez, 107
- Martínez, R., S. Mendoza, E. Reguera, P. Ortiz, and J. de la Luz Martínez. Kinetic approach to nixtamalization of corn pericarp, 107
- Mauromoustakos, A. See A. L. Dillahunty, 559
- See A. A. Perdon, 205
- McClung, A. M. See C. J. Bergman, 257
- See K. L. Bett-Garber, 551
- See E. T. Champagne, 358
- See D. S. Himmelsbach, 488
- McKenzie, K. S. See K. L. Bett-Garber, 551
- See E. T. Champagne, 358
- McKenzie, R. I. H. See A. S. Wesley, 69
- McMullen, M. S. See D. C. Doehlert, 675
- See L. A. Grant, 590
- Mehra, S. K., and S. R. Eckhoff. Influence of temper duration and weight distance on system output in the corn dry-milling process, 222
- Mehra, S. K., D. K. Gupta, P. Buriak, M. E. Tumbleson, and S. R. Eckhoff. Effect of maize tempering on throughput and product yields, 210
- Mendoza, S. See R. Martínez, 107
- Merchan, M. L. See G. M. Campbell, 272
- Mercier, M., and P. Gélinas. Effect of lipid oxidation on dough bleaching, 36
- See M. D. Pearce, 354
- See C. C. Sitakalin, 391
- Meuser, F., K. G. Busch, H. Fuhrmeister, H., and K. Rubach. Foam-forming capacity of substances present in rye, 50
- Mikola, M., O. Brinck, and B.-L. Jones. Characterization of oat endoproteases that hydrolyze oat avenins, 55
- Milliken, G. M. See M. C. Pasikatan, 730
- Milovanovic, M. See N. Mladenov, 363
- Mitchell, J. R. See A. Becker, 166
- See C. W. P. Carvalho, 342
- Mitsunaga, T. See H. Tang, 111
- Mladenov, N., N. Przulj, N. Hristov, V. Djuric, and M. Milovanovic. Cultivar-environment interactions for wheat quality traits in semiarid conditions, 363
- Moers, K. See J. A. Ingelbrecht, 721
- Moldenhauer, K. A. See K. L. Bett-Garber, 551
- See E. T. Champagne, 358
- Moreau, R. A. See V. Singh, 46, 436
- Mua, J. P. See D. Sahai, 116
- Nakamura, H. N-terminal amino acid sequence analysis of endosperm proteins in Japanese hexaploid wheat, 79
- Ng, P. K. W. See H. Köksel, 26
- See L. Lee, 447
- Noda, T., T. Tohnooka, S. Taya, and I. Suda. Relationship between physicochemical properties of starches and white salted noodle quality in Japanese wheat flours, 395
- Novaro, P., F. Colucci, G. Venora, and M. G. D'Egidio. Image analysis of whole grains: A noninvasive method to predict semolina yield in durum wheat, 217
- Obuz, E., T. J. Herald, and K. Rausch. Characterization of extruded plant protein and petroleum-based packaging sheets, 97
- Ohm, J. B. See O. K. Chung, 493
- Okot-Kotber, M., A. Liavoga, K.-J. Yong, and K. Bagorogoza. Activity and inhibition of polyphenol oxidase in extracts of bran and other milling fractions from a variety of wheat cultivars, 514
- Oksman-Caldentey, K.-M. See R.-L. Heiniö, 707
- See A. Wilhelmson, 715
- Ortiz, P. See R. Martínez, 107
- Osborne, B. G., R. Jackson, and S. R. Delwiche. Note: Rapid prediction of wheat endosperm compressive strength properties using the single-kernel characterization system, 142
- Osguthorpe, D. J. See O. Parchment, 658
- Osseyi, E. S., R. L. Wehling, and J. A. Albrecht. HPLC determination of stability and distribution of added folic acid and some endogenous folates during breadmaking, 375
- Papantoniou, E., E. W. Hammond, F. Scriven, M. H. Gordon, and J. D. Schofield. Isolation of polar lipid classes from wheat flour extracts by preparative high-performance liquid chromatography, 663
- Parchment, O., P. R. Shewry, A. S. Tatham, and D. J. Osguthorpe. Molecular modeling of an unusual spiral structure present in an elastomeric wheat seed protein, 658
- Park, J. K., S. S. Kim, and K. O. Kim. Effect of milling ratio on sensory properties of cooked rice and on physicochemical properties of milled and cooked rice, 151

- Partridge, M. A. K., A. S. Hill, M. J. Blundell, and J. H. Skerritt. Two-site sandwich ELISA for discriminating different *Gli-1* (gliadin)/*Glu-3* (LMW-glutenin subunit) alleles in hexaploid wheat, 294
- Pasikatan, M. C., E. Haque, J. L. Steele, C. K. Spillman, and G. M. Milliken. Evaluation of a near-infrared reflectance spectrometer as a granulation sensor for first-break ground wheat: Studies with six wheat classes, 730
- Payo-Rodriguez, R. See G. M. Campbell, 372
- Pearce, M. D., B. P. Marks, and J.-F. Meullenet. Effects of postharvest parameters on functional changes during rough rice storage, 354
- Pedersen, J. F. See C. Hicks, 412
- Perdon, A. A., T. J. Siebenmorgen, A. Mauromoustakos, V. K. Griffin, and E. R. Johnson. Degree of milling effects on rice pasting properties, 205
- Perera, C., Z. Lu, J. Sell, and J. Jane. Comparison of physicochemical properties and structures of sugary-2 cornstarch with normal and waxy cultivars, 249
- Peterson, D. M., A. D. Budde, C. A. Henson, and B. L. Jones. Detecting corn syrup in barley malt extracts, 349
- Phillips, D. L., J. Xing, C. K. Chong, and H. Corke. Note: Quantitative analysis of benzyl modification in waxy maize starch by Fourier Transform (FT) Raman spectroscopy, 629
- Pollak, L. M. See S. K. Singh, 315, 330, 336
- Poutanen, K. See R.-L. Heiniö, 707
- See A. Wilhelmson, 715
- Prieto-Linde, M. L. See E. Johansson, 19
- Proctor, A. See H. S. Lam, 498
- Proost, P. See H. Goesaert, 453
- Przulj, N. See N. Mladenov, 363
- Psomas, S. See M. Liakopoulou-Kyriakides, 603
- Pyun, Y. R. See J. K. Jang, 186
- Rane, K. D., and M. Cheryan. Membrane filtration of corn steep water, 400
- Rausch, K. See E. Obuz, 97
- Reguera, E. See R. Martinez, 107
- Reinikainen, P. See E.-L. Bergman, 144
- Repeckiene, A., A.-C. Eliasson, G. Juodeikiene, and E. Gunnarsson. Predicting baking performance from rheological and adhesive properties of rye meal suspensions during heating, 193
- Robertson, G. H., and T. K. Cao. Farinograph responses for wheat flour dough fortified with wheat gluten produced by cold-ethanol or water displacement of starch, 538
- Robutti, J. See K. Seetharaman, 379
- Rooney, L. W. See T. Beta, 303
- Roozeboom, K. See G. L. Brannan, 693
- Rouau, X. See J. A. Ingelbrecht, 721
- Rowe, M. See D. Sahai, 116
- Rubach, K. See F. Meuser, 50
- Sahai, D., M. O. Buendia, and D. S. Jackson. Analytical techniques for understanding nixtamalized corn flour: Particle size and functionality relationships in a masa flour sample, 14
- , J. P. Mua, I. Surjewan, M. O. Buendia, M. Rowe, and D. S. Jackson. Alkaline processing (nixtamalization) of white Mexican corn hybrids for tortilla production: Significance of corn physicochemical characteristics and process conditions, 116
- Salmenkallio-Marttila, M., K. Katina, and K. Autio. Effects of bran fermentation on quality and microstructure of high-fiber wheat bread, 429
- Sandberg, A.-S. See E.-L. Bergman, 144
- See M. C. Zghal, 1
- Savage, A. W. J. See S. Uthayakumaran, 138
- Scanlon, M. G. See M. C. Zghal, 1
- Schofield, J. D. See E. Papanthiou, 663
- Schwarzlaff, S. S., M. G. Uriyo, J. M. Johnson, W. E. Barbeau, and C. A. Griffey. Apparent dough stickiness of selected 1BL/1RS translocated soft wheat flours, 93
- Scriven, F. See E. Papanthiou, 663
- Seaborn, B. W. See O. K. Chung, 493
- Seetharaman, K., A. Tziotis, F. Borras, P. J. White, M. Ferrer, and J. Robutti. Thermal and functional characterization of starch from Argentinean corn, 379
- Seib, P. A. See G. L. Brannan, 693
- Sell, J. See C. Perera, 249
- Setser, C. S. See G. L. Brannan, 693
- Shen, M.-C. See V. M. F. Lai, 596
- Shewry, P. R. See H. A. Bloch, 74
- See O. Parchment, 658
- Siebenmorgen, T. J. See A. L. Dillahunty, 559
- See A. A. Perdon, 205
- Singh, H., and F. MacRitchie. Use of sonication to probe wheat gluten structure, 526
- Singh, S. K., L. A. Johnson, L. M. Pollak, and C. R. Hurburgh. Compositional, physical, and wet-milling properties of accessions used in Germplasm Enhancement of Maize project, 330
- , —, —, and —. Heterosis in compositional, physical, and wet-milling properties of adapted \times exotic corn crosses, 336
- , —, P. J. White, J.-L. Jane, and L. M. Pollak. Thermal properties and paste and gel behaviors of starches recovered from accessions used in the Germplasm Enhancement of Maize project, 315
- Singh, V., R. A. Moreau, and P. H. Cooke. Effect of corn milling practices on aleurone layer cells and their unique phytosterols, 436
- , —, K. B. Hicks, and S. R. Eckhoff. Effect of alternative milling techniques on the yield and composition of corn germ oil and corn fiber oil, 46
- See D. B. Johnston, 405
- Sinn, L. See A. Lapveteläinen, 322
- Sironi, E., N. Guerrieri, and P. Cerletti. Surface properties of gluten investigated by a fluorescence approach, 476
- Sitakalin, C., and J.-F. C. Meullenet. Prediction of cooked rice texture using an extrusion test in combination with partial least squares regression and artificial neural networks, 391
- See H. Köksel, 26
- Skerritt, J. H. See M. A. K. Partridge, 294
- Spillman, C. K. See M. C. Pasikatan, 730
- Stamatoudis, M. See M. Liakopoulou-Kyriakides, 603
- Steele, J. L. See M. C. Pasikatan, 730
- Steffe, J. F. See H. Köksel, 26
- See L. Lee, 447
- Stoddard, F. L. See S. Uthayakumaran, 138
- Suda, I. See T. Noda, 395
- Sun, M. See Y. Tan, 666
- Sun, X. S. See L. Carson, 101
- Sundberg, B. See L. H. Dimberg, 278
- Sunnerheim, K. See L. H. Dimberg, 278
- Suortti, T. See E. Vesterinen, 442
- See A. Wilhelmson, 715
- Surjewan, I. See D. Sahai, 116
- Swanson, B. G. See M.-R. Lee, 701
- Takeda, Y. See H. Tang, 111
- Tan, Y., Y. Xing, Q. Zhang, M. Sun, and H. Corke. Quantitative genetic basis of gelatinization temperature of rice, 666
- Tang, H., H. Ando, K. Watanabe, Y. Takeda, and T. Mitsunaga. Fine structures of amylose and amylopectin from large, medium, and small waxy barley starch granules, 111
- Tatham, A. S. See O. Parchment, 658
- See S. Uthayakumaran, 138
- Taya, S. See T. Noda, 395
- Taylor, J. R. N. See T. Beta, 303
- Thompson, D. B. See J. O. Brumovsky, 680
- Tian, Y., P. Buriak, and S. R. Eckhoff. Effect of hybrid and physical properties of individual popcorn kernels on expansion volume, 578
- Tohnooka, T. See T. Noda, 395
- Tömösközi, S. See S. Uthayakumaran, 138
- Toro-Vazquez, J. F., and C. A. Gómez-Aldapa. Chemical and physicochemical properties of maize starch after industrial nixtamalization, 543
- Tuinstra, M. R. See S. R. Bean, 84
- See C. Hicks, 412
- Tumbleson, M. E. See S. K. Mehra, 210
- Tziotis, A. See K. Seetharaman, 379
- Uriyo, M. G. See S. S. Schwarzlaff, 93
- Uthayakumaran, S., S. Tömösközi, A. S. Tatham, A. W. J. Savage, M. C. Gianibelli, F. L. Stoddard, and F. Bekes. RAPID COMMUNICATION: Effects of gliadin fractions on functional properties of wheat dough depending on molecular size and hydrophobicity, 138
- Van Damme, J. See H. Goesaert, 453
- Venora, G. See P. Novaro, 217
- Vesterinen, E., T. Suortti, and K. Autio. Effects of preparation temperature on gelation properties and molecular structure of high-amylose maize starch, 442
- Vignaux, N. See L. A. Grant, 590
- Vinyard, B. T. See R. J. Bryant, 131
- Walsh, K. See L. H. Dimberg, 278
- Wang, D., F. E. Dowell, and D. S. Chung. Assessment of heat-damaged wheat kernel using near-infrared spectroscopy, 625
- Wang, L., and Y.-J. Wang. Note: Comparison of protease digestion at neutral pH with alkaline steeping method for rice starch isolation, 690

- Wang, Y.-J. *See* L. Wang, 690
 Watanabe, K. *See* H. Tang, 111
 Wehling, R. L. *See* E. S. Osseyi, 375
 Wehrle, K. *See* S. Kenny, 458
 Wesley, A. S., O. M. Lukow, R. I. H. McKenzie, N. Ames, and D. Brown.
 Effect of multiple substitution of glutenin and gliadin proteins on flour
 quality of Canada Prairie Spring wheat, 69
 Whallon, J. H. *See* L. Lee, 447
 White, P. J. *See* K. Seetharaman, 379
 —. *See* S. K. Singh, 315
 Wieser, H. *See* S. Antes, 8, 157
 Wilhelmson, A., K.-M. Oksman-Caldentey, A. Laitila, T. Suortti, A. Kaukovirta-
 Norja, and K. Poutanen. Development of a germination process for producing
 high β -glucan, whole grain food ingredients from oat, 715
 Willett, J. L. Packing characteristics of starch granules, 64
 Wirtz, D. *See* J. Xu, 181
 Wrigley, C. W. *See* M. C. Gianibelli, 635
- Xing, J. *See* D. L. Phillips, 629
- Xing, Y. *See* Y. Tan, 666
 Xu, J., J. A. Bietz, F. C. Felker, C. J. Carriere, and D. Wirtz. Rheological
 properties of vital wheat gluten suspensions, 181
- Yeh, A.-I. *See* C.-M. Chen, 261
 —. *See* V. M. F. Lai, 596
 —. *See* Y. Lin, 647
 Yeom, H. W. *See* H. Y. L. Kim, 267
 Yong, K.-J. *See* M. Okot-Kotber, 514
- Zeller, F. J. *See* S. L. K. Hsam, 521
 Zghal, M. C., M. G. Scanlon, and H. D. Sapirstein. Effects of flour strength,
 baking absorption, and processing conditions on the structure and mech-
 anical properties of bread crumb, 1
 Zhang, Q. *See* Y. Tan, 666
 Zhu, J., and K. Khan. Effects of genotype and environment on glutenin
 polymers and breadmaking quality, 125
 —, and —. Separation and quantification of HMW glutenin subunits by
 capillary electrophoresis, 737

Subject Index

Page numbers of errata are in *italic*.

- Alkaline, steeping method; rice starch isolation, protease digestion at neutral
 pH compared (Wang and Wang), 690
 Amino acid, sequencing; of Japanese hexaploid wheat (Nakamura), 79
 α -Amylase
 —activity; prediction affected by falling number sample weight (Finney), 485
 —antistaling effect in bread and bread models (Hug-Iten et al), 421
 —synergistic hydrolysis of corn starch by (Liakopoulou-Kyriakides et al), 603
 Amylopectin
 —fine structures of, from waxy barley starch granules (Tang et al), 111
 —plasticization of side chains (Donald), 307
 Amylose
 —content; effect on expansion of extruded rice pellet (Chen and Yeh), 261
 —content of milled rice; by NIR-FT/Raman spectroscopy (Himmelsbach et
 al), 488
 —fine structures of, from waxy barley starch granules (Tang et al), 111
 —influence on wheat starch properties and breadmaking quality of starch and
 gluten blends (Lee et al), 701
 Antioxidants, avenanthramides, *p*-coumaric, ferulic and caffeic acids (Dimberg
 et al), 278
 Arabinoxylans
 —functionality; in breadmaking (Courtin et al), 564
 —influence on pasta processing and quality (Ingelbrecht et al), 721
 Artificial neural network
 —major calibration technique used for determination of moisture and protein
 in cereals (Büchmann et al), 572
 —prediction using extrusion test with partial least squares regression to pre-
 dict cooked rice texture (Sitakalin and Meullenet), 391
 Avenanthramides, stability of (Dimberg et al), 278
- Baking
 —bagels; effect of additives and storage temperature on staling properties (Lent
 and Grant), 619
 —bread; application of oat oil in (Erazo-Castrejón et al), 243
 —bread; microscale, protein additives (Antes and Wieser), 157
 —characteristics; HWW flours using computer-analyzed mixograph parameters
 (Chung et al), 493
 —micro-baking test (Hsam et al), 521
 —properties of frozen dough; influence of sodium caseinate and whey pro-
 tein (Kenny et al), 458
 —rye bread (Repeckiene et al), 193
- Barley
 —cell walls; solubilization of material from (Kanauchi and Bamforth), 121
 —malt extract; corn syrup detection in (Peterson et al), 349
 —purification and characterization of an endoxylanase inhibitor (Goesaert et
 al), 453
 —starch isolation by-products; from a starch isolation procedure (Andersson
 et al), 507
 —waxy starches; fine structures of (Tang et al), 111
- Bran
 —fermentation; effect on quality and microstructure of high-fiber wheat bread
 (Salmenkallio-Marttila et al), 429
 —fiber; cellulose/arabinoxylan mixtures from (Doner and Johnston), 200
- Bread
 —baking; application of oat oil in (Erazo-Castrejón et al), 243
 —creep-recovery of, correlation to sensory measurements of textural attributes
 (Carson and Sun), 101
 —crumb grain; effect on mechanical properties (Zghal et al), 1
 —high-fiber wheat, bran fermentation effect on quality and microstructure
 (Salmenkallio-Marttila et al), 429
 —influence of α -amylase on staling (Hug-Iten et al), 421
 —quality related to rheological measurements of wheat flour dough (Autio et al),
 654
 —retention of avenanthramides in (Dimberg et al), 278
 —surfactant effect on aeration and measurement of proofing rate (Campbell
 et al), 272
 —volume; effect of gluten proteins (Antes and Wieser), 157
- Breadmaking
 —changes in folate levels during bread production (Osseyi et al), 375
 —effect of environment on quality (Zhu and Khan), 125
 —effect of gliadin on breadmaking (Uthayakumaran et al), 138
 —functionality of arabinoxylans and endoxylanases (Courtin et al), 564
 —quality of starch and gluten blends; influence of amylose content on (Lee
 et al), 701
 —significance *Aegilops tauschii* genes (Hsam et al), 521
 —wheat cultivar and nitrogen application effects on (Johansson et al), 19
- Cake, yellow layer, shortening replacement in, by corn dextrins (Kim et al),
 267
 Cell walls, enzymolysis of (Kanauchi and Bamforth), 121
 Cellulose/arabinoxylan mixture, water binding (Doner and Johnston), 200
 Chromatography, size exclusion, reoxidized proteins (Antes and Wieser), 8
 Computer modeling, of spiral structure in elastomeric wheat seed protein
 (Parchment et al), 658
- Corn (see also Maize)
 —crosses; heterosis in compositional, physical, and wet-milling properties of
 (Singh et al), 336
 —dextrins; shortening replacement in yellow layer cakes (Kim et al), 267
 —dry-milling; influence of temper duration and weight distance on system
 output in (Mehra and Eckhoff), 222
 —fiber; cellulose/arabinoxylan mixtures from (Doner and Johnston), 200
 —germ and fiber oils; alternative milling technique effect on yield and compo-
 sition of (Singh et al), 46
 —masa flour; analytical techniques for understanding particle size and func-
 tional relationships (Sahai et al), 14
 —pericarp; corn hull, first supplement of dietary fibers in (Martínez et al), 107
 —processing; nixtamalization of white Mexican corn hybrids for tortilla pro-
 duction (Sahai et al), 116
 —steep water; membrane filtration of (Rane and Cheryan), 400
 —thermal and functional properties of starch from Argentinean corn (Seetha-
 raman et al), 379

- wet-milling; effect on aleurone layer cells and their phytosterols (Singh et al), 436
- wet-milling; proteases to reduce steep time and SO₂ requirements (Johnston and Singh), 405
- Corn starch
 - enzymic hydrolysis of, synergistic action of α -amylases and glucoamylases (Liakopoulou-Kyriakides et al), 603
 - high-amylose, preparation temperature effect on gelation and molecular structure of (Vesterinen et al), 442
 - physicochemical property and structure comparison with normal and waxy cultivars (Perera et al), 249
- Cultivar
 - by environment interactions for wheat quality traits in semiarid conditions (Mladenov et al), 363
 - normal and waxy, comparison with physicochemical properties and structures (Perera et al), 249
 - variation; effect on storage protein composition and breadmaking quality (Johansson et al), 19
 - and year effects; parameter relationships (Lapveteläinen et al), 322
- L-Cysteine, effect on rheological properties of wheat flour (Lambert and Kokini), 226
- Dehulling, of oats; optimizing conditions for (Doehlert and McMullen), 675
- Differential scanning calorimetry, to determine correlation between starch retrogradation and water mobility (Lin et al), 647
- Digestibility, rapid assay to identify high protein digestibility lines (Aboubacar et al), 160
- Dough
 - bleaching and lipid oxidation (Mercier and Gélinas), 36
 - characteristics; prediction of baking characteristics of HWW flours using computer-analyzed mixograph parameters (Chung et al), 493
 - characterization from power consumption profile of a conventional mixer (Hwang and Gunasekaran), 88
 - density measurement and proofing rates (Campbell et al), 272
 - effect of environment on properties (Zhu and Khan), 125
 - effect of heat damage on physical dough properties (Wang et al), 625
 - froze, influence of sodium caseinate and whey protein on rheology (Kenney et al), 458
 - mixing and resting; changes in SDS solubility of glutenin polymers (Aus-senac et al), 39
 - nondeveloped, partially developed, and developed, properties of (Lee et al), 447
 - rheology; effect of gluten proteins (Antes and Wieser), 157
 - wheat flour, bread quality relationship with rheological measurements of (Autio et al), 654
 - wheat flour, fortified with wheat gluten, farinograph responses for (Robertson and Cao), 538
 - wheat flour, measurement of dough stickiness in 1BL/1RS translocated soft wheat flours (Schwarzlaff et al), 93
- Drying
 - effect on sensory profile of germinated oat (Heiniö et al), 707
 - effects on functional changes during rough rice storage (Pearce et al), 354
- Dry-milling, of corn, influence of temper duration and weight distance on system output (Mehra and Eckhoff), 222
- Electrophoresis
 - capillary, for separation and quantification of HMW glutenin subunits (Zhu and Khan), 737
 - SDS-PAGE of glutenin proteins (Zhu and Khan), 125
- ELISA, for discriminating gliadin and glutenin subunit alleles in hexaploid wheat (Partridge et al), 294
- Environment, effect on glutenin protein composition (Zhu and Khan), 125
- Enzyme, inhibitor; of xylanases, purification and characterization from barley (Goesaert et al), 453
- Enzymes
 - detection of wheat preharvest sprouting using a pregelatinized starch substrate and centrifugation (Donelson et al), 282
 - transglutininase, effects of, on dough rheology (Köksel et al), 26
- Errata
 - vol. 77, no. 4, pp. 512-517, 373-374
 - vol. 77, no. 5, pp. 607-614, 105
- Extrusion
 - of maize grit, maize flour, wheat flour, and wheat semolina (Carvalho and Mitchell), 342
 - of plant protein and petroleum-based packaging sheets (Obuz et al), 97
 - of rice flour; functional and digestive characteristics (Bryant et al), 131
- Falling number
 - in rye meal suspensions (Repeckiene et al), 193
- sample weight; effects on prediction of α -amylase activity (Finney), 485
- Fatty acyl amides, starch granule accessibility to (Gray and BeMiller), 236
- Fiber
 - coarse, effect of corn milling on aleurone layer cells and their phytosterols (Singh et al), 436
 - corn, preparation of gel (Inglett and Carriere), 471
 - corn oil, alternative milling technique effect on yield and composition of (Singh et al), 46
 - resistant total dietary, of Oatrim muffins with different levels of amylose, amylopectin, and β -glucan (Li et al), 387
 - soluble, production of high-quality pasta with (Ingelbrecht et al), 721
 - wheat bran, digestibility and gastrointestinal measurements in rat (Kahlon et al), 481
 - of wheat bread; bran fermentation effect on quality and microstructure of (Salmenkallio-Marttila et al), 429
- Flavor, categorizing rice cultivars based on (Bett-Garber et al), 551
- Flour
 - benzoyl peroxide, bleaching effect of fat and lipase on carotene degradation (Mercier and Gélinas), 36
 - bug-damaged, effects of transglutininase enzyme on (Köksel et al), 26
 - corn, made from nixtamalized corn and dehydrated (Martínez et al), 107
 - nixtamalized corn, analytical techniques for understanding particle size and functionality relations (Sahai et al), 14
 - quality; effect of multiple substitution of glutenin and gliadin proteins (Wesley et al), 69
 - strength effect on dough aeration during mixing (Campbell et al), 272
 - wheat for tortillas, starch damage effects on texture (Mao and Flores), 286
 - wheat, two endoxylanases for understanding arabinoxylan functionality in breadmaking (Courtin et al), 564
- Fluorescence, approach; surface properties of gluten investigated by (Sironi et al), 476
- Foam, forming capacity of water-soluble rye proteins (Meuser et al), 50
- Folate, changes in native folate levels during breadmaking (Osseyi et al), 375
- Folic acid, changes in concentration of added, during breadmaking (Osseyi et al), 375
- FT-Raman spectroscopy, benzylated starch analysis (Phillips et al), 629
- Functional properties, of starch from Argentinean corn (Seetharaman et al), 379
- Functionality, changes during rough rice storage (Pearce et al), 354
- Gelatinization, temperature of rice; quantitative genetic basis of (Tan et al), 666
- Germ, corn oil, alternative milling technique effect on yield and composition of (Singh et al), 46
- Glass transition, temperature
 - of maize and wheat extrudates at various water contents by DMTA (Carvalho and Mitchell), 342
 - T_g of starch (Toto-Vazquez and Gómez-Aldapa), 543
- Gliadin
 - contribution to functional properties of wheat flour dough (Uthayakumaran et al), 138
 - ELISA for discriminating different alleles in hexaploid wheat (Partridge et al), 294
 - protein; multiple substitution effect of, on Canada Prairie Spring wheat flour quality (Wesley et al), 69
- β -Glucan
 - germination process for producing; whole grain food ingredients from oat (Wilhelmson et al), 715
 - solubilization of (Kanauchi and Bamforth), 121
- Glutelins, reoxidation of subunits (Antes and Wieser), 8
- Gluten
 - proteins; sonication to probe structure (Singh and MacRitchie), 526
 - proteins; wheat gliadin and glutenin proteins (Bean and Lookhart), 608
 - and starch blends, influence of amylose content on wheat starch properties and breadmaking qualities of (Lee et al), 701
 - surface properties of, fluorescence approach (Sironi et al), 476
 - wheat, farinograph responses for wheat flour dough fortified with (Brites and Carrillo), 59
- Glutenin
 - effect of environment on composition (Zhu and Khan), 125
 - HMW and LMW subunits, reoxidation, effect on dough rheology and bread volume (Antes and Wieser), 157
 - polymers; changes in SDS solubility of, during dough mixing and resting (Aus-senac et al), 39
 - protein; multiple substitution effect of, on Canada Prairie Spring wheat flour quality (Wesley et al), 69
 - sonication to probe structure (Singh and MacRitchie), 526
 - subunits; ELISA for discriminating different alleles in hexaploid wheat (Partridge et al), 294
 - subunits of wheat, preparative method for in vitro production of functional polymers from (Beasley et al), 464

- wheat, biochemical, genetic, and molecular characterization of (Gianibelli et al), 635
- Grain sorghum, hybrids; sensory characteristics of, for potential human food use (Brannan et al), 693
- High molecular weight, and LMW glutenin subunits; controlled by *Glu-1* and *Glu-3* loci on durum wheat quality (Brites and Carrillo), 59
- High-amylose, maize starch; preparation temperature effect on gelation and molecular structure of (Vesterinen et al), 442
- High-performance capillary electrophoresis, method using a narrow bore glass capillary and an electric field to separate components in solution (Bean and Lookhart), 530
- HMW, glutenin subunits; separation and quantification by capillary electrophoresis (Zhu and Khan), 737
- HPLC
 - determination of folates in bread (Osseyi et al), 375
 - isolation of polar lipid classes from wheat flour extracts by (Papantoniou et al), 663
- Hybrid, of popcorn, effect on expansion volume (Tian et al), 578
- Hydrothermal processing, to degrade phytate; heat and moisture for (Bergman et al), 144
- Hydrothermal treatment, of high-amylose maize starch; boiling-stable granular resistant starch produced by (Brumovsky and Thompson), 680
- Image analysis
 - of crumb grain; quantification of crumb structure (Zghal et al), 1
 - noninvasive technique for studying the relationships among grain size and shape descriptors and semolina yield (Novaro et al), 217
- Instructions to authors, iv
- Lipid
 - classes, isolation of from wheat flour extracts by preparative HPLC (Papantoniou et al), 663
 - oxidation and dough bleaching (Mercier and Gélinas), 36
- Maize (see also Corn)
 - high-amylose starch, partial acid hydrolysis and hydrothermal treatments of, for production of boiling-stable granular resistant starch (Brumovsky and Thompson), 680
 - hulls; cellulosic fiber gels prepared from cell walls of (Inglett and Carriere), 471
 - project; Germplasm Enhancement of Maize, wet-milling properties used in (Singh et al), 330
 - proteins extracted by SDS and separated by capillary electrophoresis (Bean et al), 84
 - starch; chemical and physicochemical properties of after nixtamalization (Toto-Vazquez and Gómez-Aldapa), 543
 - tempering effect on throughput and product yields (Mehra et al), 210
 - thermal and pasting properties and starch gel recovered from accessions in germplasm enhancement of (Singh et al), 315
 - zein protein functionality (Bugusu et al), 31
- Malt, barley extract, dilution with corn syrup (Peterson et al), 349
- Methods
 - capillary electrophoresis, separation of sugars (Doner and Johnston), 200
 - microsatellite in rice waxy gene to determine amylose class (Bergman et al), 257
 - preparative, for in vitro production of functional polymers from (Beasley et al), 464
 - rapid, for milled rice surface total lipids and free fatty acids (Lam and Proctor), 498
- Microscopy, using laser scanning confocal microscopy to examine protein network in dough (Lee et al), 447
- Milling
 - degree of effects on rice pasting properties (Perdon et al), 205
 - effect of on RVA profile of starch products (Becker et al), 166
 - fractions; activity and inhibition of polyphenol oxidase in bran and shorts (Okot-Kotber et al), 514
- Mixing, characterization of wheat dough using a conventional mixer (Hwang and Gunasekaran), 88
- Mixograph, prediction of baking characteristics, continuum regression (Chung et al), 493
- Moisture
 - parameter analyzed in wheat and barley (Büchmann et al), 572
- Multangle laser light scattering, technique used to determine absolute molecular weight of components in solution by measuring laser light scattering at multiple angles from incident laser (Bean and Lookhart), 608
- Near-infrared reflectance, analysis for prediction of cooked rice texture (Cham-pagne et al), 358
- Near-infrared transmission, analytical technique used for determination of
 - moisture and protein in cereals (Büchmann et al), 572
- Near-isogenic wheat lines, to determine protein composition-functionality relationships (MacRitchie and Lafandra), 501
- NIR spectrometer, as granulation sensor for first-break ground wheat (Pasikatan et al), 730
- NIR spectroscopy
 - classification of heat-damaged wheat (Wang et al), 625
- Nitrogen, application; effect on storage protein composition and breadmaking quality (Johansson et al), 19
- Nixtamalization
 - alkali-temperature process for maize (Toto-Vazquez and Gómez-Aldapa), 543
 - of white Mexican corn hybrids for tortilla production (Sahai et al), 116
 - process to make tortillas in which whole corn is cooked in lime water and steeped (Martínez et al), 107
- NMR, to determine correlation between starch retrogradation and water mobility (Lin et al), 647
- Noodles
 - effect of starch properties on white salted quality, relationships in Japanese wheat flours (Noda et al), 395
 - sorghum starch (Beta and Corke), 417
- Oat
 - antioxidants (Dimberg et al), 278
 - dehulling; optimizing conditions for (Doehlert and McMullen), 675
 - effect on sensory profile of germinated (Heiniö et al), 707
 - extract containing high levels of β -glucan (Li et al), 387
 - germination process for producing high β -glucan, whole grain food ingredients from (Wilhelmsen et al), 715
 - oil, application of in breadbaking (Erazo-Castrejón et al), 243
 - physical, chemical, and sensory parameters of grains, groats, rolled oats and oatmeal; cultivar and year effects (Lapveteläinen et al), 322
 - proteinases of germinated avenin hydrolyzing enzymes (Mikola et al), 55
- Oxidation, glutenin subunits KBrO₃ and KIO₃ (Antes and Wieser), 8
- Packing fraction, of granules from various starches (Willett), 64
- Pasta, processing; influence of arabinoxylans and endoxylanases on (Ingelbrecht et al), 721
- Physical properties, of popcorn, effect on expansion volume (Tian et al), 578
- Phytate, antinutrient that chelates minerals (Bergman et al), 144
- Polyphenol oxidase, discoloring enzyme in wheat-based foods (Okot-Kotber et al), 514
- Popcorn
 - expansion volume affected by hybrid and physical properties of (Tian et al), 578
- Power consumption, used to determine dough mixing characteristics (Hwang and Gunasekaran), 88
- Protease
 - digestion at neutral pH; comparison with alkaline steeping method for rice starch isolation (Wang and Wang), 690
 - to reduce steep time and SO₂ requirements during corn wet-milling (Johnston and Singh), 405
- Protein
 - categorizing rice cultivars based on (Bett-Garber et al), 551
 - composition of storage proteins in sorghum (Hicks et al), 412
 - composition-functionality in near-isogenic wheat lines (MacRitchie and Lafandra), 501
 - content of milled rice; by NIR-FT/Raman spectroscopy (Himmelsbach et al), 488
 - effect of drying on protein determination (Wang et al), 625
 - extraction of maize and sorghum proteins with SDS (Bean et al), 84
 - formation of protein matrix during dough development (Lee et al), 447
 - parameter analyzed in wheat and barley (Büchmann et al), 572
 - wheat seed, spiral structure present in, molecular modeling of (Parchment et al), 658
- Proteinases, of germinated oats; which hydrolyze avenins and oat prolamins (Mikola et al), 55
- Rheological properties
 - of high-amylose maize starch, preparation temperature effect (Vesterinen et al), 442
 - of maize fiber gel (Inglett and Carriere), 471
- Rheology
 - of crumb grain; relationship to structure (Zghal et al), 1
 - of frozen dough; influence of sodium caseinate and whey protein (Kenny et al), 458
 - of gluten from diploid, tetraploid, and hexaploid wheat (Hsam et al), 521
 - relationships between rheological properties and microstructural characteristics of doughs (Lee et al), 447
 - rye meal suspensions (Repeckiene et al), 193

- studies of bug-damaged dough, effect of transglutininase enzyme on dough properties (Köksel et al), 26
- wheat dough, extensigram (Antes and Wieser), 157
- of wheat gluten suspensions (Xu et al), 181
- Rice
 - cultivars; categorizing (Bett-Garber et al), 551
 - degree of milling; sensory properties of cooked rice and physicochemical properties of milled and cooked rice (Park et al), 151
 - expansion of extruded pellet (Chen and Yeh), 261
 - extruded flour, functional and digestive characteristics (Bryant et al), 131
 - functional changes during storage (Pearce et al), 354
 - gelatinization temperature of, quantitative genetic basis of (Tan et al), 666
 - method for amylose class (Bergman et al), 257
 - milled, protein and apparent amylose contents of, by NIR-FT/Raman spectroscopy (Himmelsbach et al), 488
 - milled, rapid methods for surface total lipids and free fatty acids (Lam and Proctor), 498
 - pasting properties; effect of degree of milling (Perdon et al), 205
 - starch isolation; alkaline steeping method for, protease digestion at neutral pH compared (Wang and Wang), 690
 - starch, molecular and gelatinization properties (Lai et al), 596
 - texture, cooked; NIR analysis for prediction of (Champagne et al), 358
 - texture; prediction using extrusion test with partial least squares regression and artificial neural network (Sitakalin and Meullenet), 391
- Rye
 - HMW glutelin subunits, reoxidation (Antes and Wieser), 8
 - meal suspension; rheology of (Repeckiene et al), 193
 - predictive models for optimization of phytate degradation (Bergman et al), 144
 - water-soluble proteins, foam-forming capacity of (Meuser et al), 50
- SCIRA, detecting corn syrup on barley malt extract (Peterson et al), 349
- SDS, use of to extract maize and sorghum proteins (Bean et al), 84
- Self-assembly, starch granule structure (Donald), 307
- Semolina
 - quality; break-roll differential effect on quality (Manthey and Hareland), 368
 - yield, a noninvasive method to predict (Novaro et al), 217
- Shortening, replacer; by corn dextrins in yellow layer cakes (Kim et al), 267
- Size-exclusion chromatography, separation of components by size using chromatographic supports (Bean and Lookhart), 608
- Sonication, of wheat gluten to probe structure (Singh and MacRitchie), 526
- Sorghum
 - analysis of kafirin composition and feed quality traits (Hicks et al), 412
 - composite flour bread (Bugusu et al), 31
 - high protein digestibility mutant (Aboubacar et al), 160
 - proteins extracted by SDS and separated by capillary electrophoresis (Bean et al), 84
 - starch; genetic diversity (Beta et al), 583
 - starch, steeping treatment effect on pasting and thermal properties of (Beta et al), 303
 - starch noodles (Beta and Corke), 417
- Spaghetti, quality; break-roll differential effect on quality (Manthey and Hareland), 368
- Staling, bagels; effect of additives and storage temperature on (Lent and Grant), 619
- Starch
 - benzylated, FT-Raman analysis (Phillips et al), 629
 - boiling-stable granular resistant, produced by partial acid hydrolysis and hydrothermal treatments of maize starch (Brumovsky and Thompson), 680
 - and by-products; from a barley starch isolation procedure (Andersson et al), 507
 - conversion; of maize and wheat extrudates by DSC, X-ray, and Rapid Visco Analyser (Carvalho and Mitchell), 342
 - damage; effect on wheat flour tortilla texture (Mao and Flores), 286
 - degradation by α -amylase in bread (Hug-Iten et al), 421
 - detection of wheat preharvest sprouting using a pregelatinized starch substrate and centrifugation (Donelson et al), 282
 - evaluation of conversion using DSC, alkaline viscosity, and RVA (Becker et al), 166
 - gel, recovered from accessions in germplasm enhancement of maize, thermal and pasting properties (Singh et al), 315
 - gelatinization; swelling of starch granules and melting of starch structure (Toto-Vazquez and Gómez-Aldapa), 543
 - and gluten blends; influence of amylose content on wheat starch properties and breadmaking quality of (Lee et al), 701
 - granule; plasticization and self-assembly in (Donald), 307
 - granule reaction sites (Huber and BeMiller), 173
 - granules; accessibility of (Gray and BeMiller), 236
 - granules; packing characteristics (Willett), 64
 - method for rice amylose class (Bergman et al), 257

- noodle quality (Beta and Corke), 417
- protein interactions (Bloch et al), 74
- relation to white salted noodle quality in Japanese wheat flours (Noda et al), 395
- resistant, of Oatrim muffins with different levels of amylose, amylopectin, and β -glucan (Li et al), 387
- retrogradation; water mobility correlation, determined by DSC and NMR (Lin et al), 647
- rice, molecular and gelatinization properties, effect of amylose content (Lai et al), 596
- sorghum, genetic diversity (Beta et al), 583
- sorghum, steeping treatment effect on pasting and thermal properties of (Beta et al), 303
- thermal and functional properties from Argentinean corn (Seetharaman et al), 379
- waxy and nonwaxy durum wheats, characteristics of (Grant et al), 590
- wheat, effect of glass transition, gelatinization, and retrogradation (Jang et al), 186
- Steep water, corn, membrane filtration of (Rane and Cheryan), 400
- Steeping, time; proteases to reduce (Johnston and Singh), 405
- Storage
 - protein composition; wheat cultivar and nitrogen application effects on (Johansson et al), 19
 - rough rice, effects of postharvest parameters on functional changes (Pearce et al), 354
- Sugar, effect on expansion indices, specific mechanical energy, glass transition temperature, and starch conversion of maize and wheat extrudates (Carvalho and Mitchell), 342
- Surfactants, effect on dough aeration and proofing rate (Campbell et al), 272
- Temperature
 - effect on rice color and viscosity (Dillahunty et al), 559
 - gelatinization, rice, quantitative genetic basis of (Tan et al), 666
- Tempering
 - maize, effect on throughput and product yields (Mehra et al), 210
 - process of adding water to corn kernel to make pliable embryo (Mehra and Eckhoff), 222
- Texture
 - categorizing rice cultivars based on (Bett-Garber et al), 551
 - cooked rice, NIR analysis for prediction of (Champagne et al), 358
 - cooked rice; prediction using extrusion test with partial least squares regression and artificial neural network (Sitakalin and Meullenet), 391
 - creep-recovery of bread, correlation to sensory measurements of (Carson and Sun), 101
 - wheat flour tortillas, starch damage effects on (Mao and Flores), 286
- Thermal properties
 - of starch from Argentinean corn (Seetharaman et al), 379
 - of starches recovered from accessions in germplasm enhancement of maize (Singh et al), 315
- Tortillas
 - flat and rounded cooked cakes made from nixtamalized corn (Martínez et al), 107
 - nixtamalization of white Mexican corn hybrids for production of (Sahai et al), 116
 - wheat flour, starch damage effects on texture (Mao and Flores), 286
- Viscosity
 - effect of milling on apparent viscosity as shown by RVA (Becker et al), 166
 - of maize fiber gel (Inglett and Carriere), 471
 - measurement of starch (Dillahunty et al), 559
- Water mobility, starch retrogradation correlation, determined by DSC and NMR (Lin et al), 647
- Wet-milling
 - corn; effect on aleurone layer cells and their phytosterols (Singh et al), 436
 - corn, proteases to reduce steep time and SO_2 requirements (Johnston and Singh), 405
 - effect on yield and composition of corn germ oil and corn fiber oil (Singh et al), 46
 - properties; of accessions used in Germplasm Enhancement of Maize project (Singh et al), 330
 - properties; heterosis in, of corn crosses (Singh et al), 336
- Wheat
 - bread; bran fermentation effect on quality and microstructure (Salmenkallio-Marttila et al), 429
 - detection of wheat preharvest sprouting using a pregelatinized starch substrate and centrifugation (Donelson et al), 282
 - for discriminating gliadin and glutenin subunit alleles in hexaploid (Partridge et al), 294

- dough stickiness in 1BL/1RS and non-1BL/1RS soft wheat flours (Schwarzlaff et al), 93
- durum, break-roll differential effect on semolina and spaghetti quality (Manthey and Hareland), 368
- durum, HMW and LMW glutenin subunits controlled by *Glu-1* and *Glu-3* loci on (Brites and Carrillo), 59
- durum, a noninvasive method to predict semolina yield in (Novaro et al), 217
- durum and bread, surface properties of gluten, fluorescence approach (Sironi et al), 476
- durum starch, characteristics of waxy and nonwaxy (Grant et al), 590
- endosperm compressive strength; rapid prediction (Osbourne et al), 142
- enzymes; activity and inhibition of polyphenol oxidase in bran and shorts (Okot-Kotber et al), 514
- falling number sample weight effects on prediction of α -amylase activity (Finney), 485
- first-break ground, NIR spectrometer as granulation sensor for (Pasikatan et al), 730
- flour; effect of L-cysteine on rheological properties of (Lambert and Kokini), 226
- flour; effect of starch properties on white salted noodle quality in (Noda et al), 395
- flour dough; bread quality relationship with rheological measurements of (Autio et al), 654
- flour dough; fortified with wheat gluten, farinograph responses for (Robertson and Cao), 538
- flour extracts; isolation of polar lipid classes from, by preparative HPLC (Papantoniou et al), 663
- glutenin; biochemical, genetic, and molecular characterization of (Gianibelli et al), 635
- glutenin subunits of, preparative method for in vitro production of functional polymers from (Beasley et al), 464
- HMW and LMW subunits, reoxidation (Antes and Wieser), 8
- predictive models for optimization of phytate degradation in (Bergman et al), 144
- quality traits; cultivar-by-environment interactions for, in semiarid conditions (Mladenov et al), 363
- starch; effect of glass transition, gelatinization, and retrogradation (Jang et al), 186
- starch; influence of amylose content on (Lee et al), 701
- use of near-isogenic wheat lines to determine composition-functionality relationships (MacRitchie and Lafiandra), 501
- Wheat bran
 - fiber; digestibility and gastrointestinal measurements in rat (Kahlon et al), 481
 - mechanical properties (Mabille et al), 231
- Xylanase, inhibitor; purification and characterization from barley (Goesaert et al), 453
- Yellowing, quality defect and discoloration of rice grains (Dillahunty et al), 559
- Zein, viscoelastic fibril formation (Bugusu et al), 31